

Version 3, 2nd April 2020

iADH COVID-19 Factsheet

*Disclaimer: The aim of this factsheet is to provide an easily consultable list of facts regarding the current COVID-19 pandemic for iADH members. These facts will evolve and change as knowledge grows and therefore cannot be considered definitive or in any way binding. Scientific publications have been sought whenever available, but some information presented here has been sourced from different government or professional guidelines and recommendations. This factsheet is not a substitute for professional medical advice. **All members must follow their national guidelines as situations will differ greatly between countries and regions.***

The virus

- COVID-19 is the infectious disease caused by the most recently discovered coronavirus. This new virus and disease were unknown before the outbreak began in Wuhan, China, in December 2019 (1).
- The new virus is called SARS-CoV-2 (1).
- No medical treatment or vaccine has as yet been found for COVID-19.
- The virus is more contagious than seasonal flu (1).
- The cell receptor for SARS-CoV-2 is angiotensin-converting enzyme II (ACE2), a receptor that is highly expressed on the oral mucosa and tongue (2)

Incubation period and recovery period

- The time between infection and experiencing symptoms is estimated at an average of 5 days, with a usual range of 3 to 7 days, and a maximum of 14 days (1).
- Mild cases of COVID-19 have an early viral clearance with 90 percent of patients repeatedly testing negative at 10 days post-onset, by contrast persons with severe COVID-19 have a high viral load and a long virus-shedding period (remain positive at or beyond 10 days post-onset) (3)

Symptoms

- The most common symptoms of COVID-19 are fever, dry cough, tiredness/fatigue and dyspnoea/shortness of breath (4). Some patients may have muscular aches and pains, headache, nasal congestion, runny nose, sore throat, confusion, vomiting, diarrhoea or anosmia (5). These symptoms begin gradually (1, 6).
- Some infected persons will have no symptoms or very mild symptoms but are still contagious (7).
- Around 15% (1 in 6) of infected people will develop severe symptoms, such as acute respiratory distress syndrome, arrhythmia and shock (6).
- Of the 15% with severe disease, it has been estimated that 5% will need intensive care facilities. The mortality rate in intensive care is between 40 and 70% (8).

Patient groups

- Persons of all ages are susceptible to COVID-19 (6), even in its most severe forms.
- Children are more likely to have asymptomatic forms
- Groups more likely to develop severe symptoms include (6,9):
 - Persons over 70 years of age
 - History of cardiovascular disease: severe hypertension, stroke, cardiac surgery, cardiac arrest, cardiac failure
 - Unstable insulin-dependent diabetes
 - Chronic respiratory disease
 - Renal failure and dialysis
 - Cancer patients currently undergoing treatment
 - Congenital or acquired immune deficiency disorders
 - Liver cirrhosis
 - Morbid obesity
 - Pregnant women
- It may be useful to consider the entire population as falling into one of four groups:
 - Confirmed cases – positive testing for virus and/or positive CT imaging (these may not be available for many cases or in some countries)
 - Suspected cases – people with an epidemiological history, COVID-19 related symptoms and / or positive CT imaging
 - Recovered cases – sometimes described as two consecutive negative nasopharyngeal swab specimens collected \geq 24hours apart (10)
 - At risk population – everybody not in one of the other groups
- Dentists are advised to take the temperature of all patients on arrival at the surgery (11)

Diagnosis

- The reliability of available tests is unknown but the literature reports that a single negative RT-PCR test result from suspected patients does not exclude infection (6).
- In severe cases of COVID-19 chest imaging can show bilateral pneumonia (6).

At-risk professionals

- Dentists are amongst those health professionals most at risk (12)
- Those who are in close contact with symptomatic and asymptomatic COVID-19 patients, such as healthcare workers and other patients in hospital are at risk (6).

At the Wuhan University Hospital dental school, using adequate protection measures (see dental treatment section below), 169 staff were involved in the emergency treatment of > 700 patients. Nine staff members contracted COVID-19 (6).

Transmission

- Asymptomatic people and people during the incubation period are also carriers of SARS-CoV-2 (6,7).
- The virus is most frequently spread by droplets (>5µm diameter) and aerosol (<5µm diameter) from the nose or mouth, on talking, coughing, sneezing, laughing etc. (1,2,13).
- Infection can occur through mucosa and through ocular contamination (14)
- Droplets landing on objects or surfaces remain contagious for up to 3 hours on dry surfaces and up to 9 days on damp surfaces (15).
- If a person touches a contaminated object/surface and then touches their face (eyes, nose or mouth) they may become infected (1).
- Transformation of saliva, blood and mucosal secretions into an aerosol increases the risk of transmission (e.g when using high-speed dental handpiece or ultrasonic instruments) (2,16)

Prevention through telephone triage

- Patients should be encouraged to phone services prior to coming into emergency services without an appointment (17)
- Where possible treatment of a known COVID 19 patient should be deferred for at least one month (1)
- Emergency treatment for recovered patients is possible at least 3 days following recovery and at least 7 days after first symptoms (17).
- Emergency treatment for asymptomatic confirmed cases is possible at least 7 days after diagnosis if there has been no subsequent illness (17).
- Dental treatment should be avoided whenever possible (6,17) and telephone triage should aim to provide advice and access to prescription rather than treatment when possible.
- Questions aiming to screen for COVID-19 include (2,11):
 - COVID-19 suspected or confirmed?
 - Fever over the last 14 days?
 - Respiratory problems, cough, breathlessness over the last 14 days?
 - Direct contact with a confirmed or suspected patient?

Preventing droplet transmission

- A distance of at least 1 metre (3 feet), preferably 1.5-2 metres (6 feet), should be maintained between people so that if they cough or sneeze they do not contaminate each other (1,16)
 - Cover your mouth and nose with a tissue or your bent elbow when you cough or sneeze. Dispose of the used tissue immediately into a covered bin. (1)
- Avoidance of use of high-speed dental handpieces or ultrasonic instruments (16).

Preventing contact transmission from objects/ surfaces

- Surfaces can be disinfected using 62-71% ethanol; or 0.5% hydrogen peroxide; or 0.1% sodium hypochloride with surface contact time of one minute. (18)
- 0.1% sodium hypochloride can be prepared from household bleach at 2.6% concentration. Mix 100 ml (millilitres) of 2.6% bleach with 2.4 litres of water to obtain 2.5 litres of solution. This solution is stable 24 hours. (18)
- Aerosol in the dental surgery contaminates the dental surgery (chair, surfaces, floor...) so careful decontamination and airing of the surgery is necessary between patients (18).
- In experimental conditions, the virus remains viable in aerosol form for over 3 hours. Following aerosol, the virus was stable on plastic and stainless steel surfaces up to 72 hours. The half-life of the virus in aerosol was 1.1 to 1.2 hours (19).
- Cleaning and decontamination immediately after a procedure may be ineffective due to later settling of airborne small particles (19) but there is no consensus on the time necessary to ensure effectiveness of cleaning in a well ventilated surgery.

Hand hygiene

- Hand hygiene is the most critical measure for reducing the risk of transmitting microorganism to the patient. (6)
- Soap and water are effective in destroying the virus (but do not remove bacterial contamination) (20)
- Alcohol-based hand disinfectant is effective in destroying the virus and removing bacterial contamination (20)

Dental treatment

- Routine dental treatment should be avoided if at all possible (6,17)
- Pre procedural mouthwash with 1% hydrogen peroxide or 0.2% povidone has been recommended. Chlorhexidine gluconate has been reported to be ineffective (11)
- Use of rotary instruments creating aerosol should be avoided whenever possible (6)
- Use of the 3-way syringe should be avoided whenever possible (6)
- Extra-oral radiography should be preferred to intra-oral radiography to reduce the risk of provoking salivation and coughing (6)
- If needed use anti-refraction high speed dental handpiece to reduce dental waterline contamination (11)
- Treatment should be performed under rubber dam to reduce contact with saliva (11)
- High volume suction should be used during treatment (6)
- Four-handed dentistry should be practised (6)
- The surgery should be well ventilated and then fully decontaminated (all surfaces and objects not forgetting door handles, chairs desks) between patients (6,11).
- If necessary, absorbable sutures are to be preferred (11)

Personal protective equipment (PPE)

- Gloves, goggles or face shield, mask, surgical uniform and gown are necessary during treatment (6) of any suspected COVID-19 patient (in many regions ALL patients may be considered as potentially infected) and dentists are also potential asymptomatic vectors. Caps and overshoes are also recommended (6,11).
- Respirator masks i.e. N-95 masks (National Institute of Occupational Safety and Health) or FFP2 or FFP3 masks (European Union standard Filtering Face Piece), are recommended for routine dental practice (16) of any suspected or confirmed COVID-19 patient (in some regions ALL patients may be considered as potentially infected)

Organisation of the dental clinic

- Unless all patients are considered as being potentially infected, confirmed COVID-19 patients should enter and wait in a separate area and be given surgical face masks (15). Separate toilet facilities should also be provided.
- If absolutely necessary to treat a confirmed COVID-19 patient because they require urgent dental care which cannot be postponed (21), they should be planned as the last patient in the day in order to be able to deep clean and ventilate the surgery. Cleaning may be repeated in the morning in case aerosol has deposited overnight.

Self-monitoring for healthcare professional

- Healthcare professionals in contact with patients should monitor themselves for fever by taking their temperature twice a day and remain alert for respiratory symptoms (e.g., cough, shortness of breath, sore throat) (22)
- Healthcare professionals should stay at home if they have any COVID-19 symptoms (17)
- Anyone self-monitoring should be prepare a plan for whom to contact if they develop fever or respiratory symptoms during the self-monitoring period to determine whether medical evaluation is needed. (22)
- Healthcare professionals with patient contact at work should follow guidance to reduce transmission in their homes (23)

Return to work after COVID-19

- Healthcare professionals should follow personalised medical advice as to their ability to return to work
- Some regions or countries prohibit return to work until at least 3 days (72 hours) have passed since recovery (resolution of fever without the use of fever-reducing medications and improvement in respiratory symptoms) and at least 7 days after any symptoms first appeared (10)
- Other countries or regions rely on testing and advice waiting for two consecutive negative nasopharyngeal swab specimens collected ≥ 24 hours apart (10)
- On return to work, it has been suggested that healthcare professionals should wear a facemask for 14 days and contact with severely immunocompromised patients (e.g., transplant, haematology-oncology) may be restricted until 14 days after illness onset (10)

People with long-term conditions and national emergencies

- People with long-term conditions are at risk of being neglected during national emergencies (24)
- Risk is described in relation to disruption of health care resources; interruption to routine care; interruption to medication supply; increased stress; changes in food supply; changes in activity level; disruptions in transport (24)
- People living in deprived areas are at increased risk and there is interaction between long term conditions and deprivation level (24)

Changes in Oral Pathology during COVID19 pandemic

- Global health emergencies have a psychological impact on the general public and can lead to changes in the incidence of different oral diseases (25). The incidence of idiopathic and/or stress-related symptoms may increase, such as recurrent aphthous ulcer, oral lichen planus, burning mouth syndrome, temporomandibular joint disorder, bruxism, atypical odontalgia etc.
- Online consultations can be considered for these patients, to help reduce the anxiety and feeling of helplessness (25).



iADH Covid-19 Fact Sheet by IADH is licensed under a

[Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-nc-sa/4.0/).

Authors: Denise Faulks and Alison Dougall, in collaboration with Suzanne Krämer, Blanaid Daly and Pedro Diz Dios.

Comments, corrections, additions are welcome: vicechair@iadh.org

REFERENCES:

- 1) WHO Q&A on coronaviruses (COVID-19). Available at: <https://www.who.int/news-room/q-a-detail/q-a-coronaviruses> Consulted 27/03/2020.
- 2) Ge ZY, Yang LM, Xia JJ, Fu XH, Zhang YZ. Possible aerosol transmission of COVID-19 and special precautions in dentistry. Journal of Zhejiang University Mar 2020 Available at <https://link.springer.com/content/pdf/10.1631/jzus.B2010010.pdf>
- 3) Liu Y, Yan LM, Wan L, Xiang TX, Le A, Liu JM, Peiris M, Poon LLM, Zhang W. Viral dynamics in mild and severe cases of COVID-19. The Lancet March 19 2020 [https://www.thelancet.com/action/showPdf?pii=S1473-3099\(20\)30232-2](https://www.thelancet.com/action/showPdf?pii=S1473-3099(20)30232-2)
- 4) Chaoqun Ma, Jiawei Gu, Pan Hou, Liang Zhang, Yuan Bai, Zhifu Guo, Hong Wu, Bili Zhang, Pan Li, Xianxian Zhao. Incidence, clinical characteristics and prognostic factor of patients with COVID-19: a systematic review and meta-analysis. medRxiv 2020.03.17. Available at <https://www.medrxiv.org/content/10.1101/2020.03.17.20037572v1>
- 5) Lüers JC, Klußmann JP, Guntinas-Lichius O. [The Covid-19 pandemic and otolaryngology: What it comes down to?] Laryngorhinootologie. 2020 Mar 26. Article in German. doi: 10.1055/a-1095-2344. [Epub ahead of print]
- 6) Meng L, Hua F & Bian Z. Coronavirus disease 2019 (COVID-2019): Emerging and future challenges for dental and oral medicine. Journal of Dental Research 2020 <https://doi.org/10.1177/0022034520914246>
- 7) Rothe C, Shunk M, Sothmann P et al. Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. N Engl J Med. 2020; 382:970-971. Doi:10.1056/NEJMc2001468
- 8) Ministère des solidarités et de la santé (France). Message d’alerte rapide sanitaire. Besoins en reanimation et réponse capacitaire. 20 mars 2020.
- 9) Haut Conseil de la santé publique (France). Avis relative à la prise en charge des patients à risqué de forme sévère de COVID-19. 14th March 2020.
- 10) Centers for Disease Control and Prevention. Coronavirus Disease 2019. Criteria for return to work for healthcare personnel with confirmed or suspected COVID-19 (interim guidance). <https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/hcp-return-work.html>
- 11) Peng X, Xu X, Li Y et al. Transmission routes of 2019-nCov and controls in dental practice. Int J Oral Sci 2020; 12,9 <https://doi.org/10.1038/s41368-020-0075-9>
- 12) Gamio L. The workers who face the greatest coronavirus risk. The New York Times, 15th March 2020. Available at: <https://www.nytimes.com/interactive/2020/03/15/business/economy/coronavirus-worker-risk.html>
- 13) Bourouiba L. Turbulent gas clouds and respiratory pathogen emissions – potential implications for reducing transmission of COVID-19. Journal of the American Medical Association March 26 2020. <https://jamanetwork.com/journals/jama/fullarticle/2763852>
- 14) To KK, Tsang OT, Chik-Yan Yip C et al. Consistent detection of 2019 novel coronavirus in saliva. Clin Infect Dis 2020 pii: ciaa149, in press
- 15) Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. J Hosp Infect 2020; 104:246-251.
- 16) Wei J, Li Y. Airborne spread of infectious agents in the indoor environment. Am J Infect Control 2016,44:S102-S108.
- 17) Centers for Disease Control and Prevention. Coronavirus Disease 2019. Interim infection, prevention and control guidance for dental settings during the COVID-19 response. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/dental-settings.html>
- 18) Haut Conseil de la santé publique (France). Avis relative au traitement du linge, au nettoyage, d’un logement ou de la chambre d’hospitalisation d’un patient confirmé à SARS-CoV-2 et à la protection des personnes. 18 February 2020.
- 19) Van Doremalen N, Bushmaker T, Morris D et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. New England Medical Journal, march 17 2020. Doi:10.1056/NEJMc2004973
- 20) Fermer R. Hand disinfectant and COVID-19. Centre for Evidence Based Medicine Oxford. March 17th 2020. Available at: <https://www.cebm.net/covid-19/hand-disinfectant-and-covid-19/>

- 21) SDCEP Management of Acute Dental Problems During COVID-19 Pandemic. 30 March 2020. Available at <http://www.sdcep.org.uk/wp-content/uploads/2020/03/SDCEP-MADP-COVID-19-guide-300320.pdf>
- 22) Centers for Disease Control and Prevention. Coronavirus Disease 2019. Interim US guidance for risk assessment and public health management of healthcare personnel with potential exposure in a healthcare setting to patients with coronavirus disease (COVID-19). <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-risk-assessment-hcp.html>
- 23) Centers for Disease Control and Prevention. Coronavirus Disease 2019. Cleaning and disinfecting your home. <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/disinfecting-your-home.html>
- 24) Hartmann-Boyce J, Mahtani KR. Supporting people with long-term conditions (LTCs) during national emergencies, Centre for Evidence Based Medicine Oxford. March 25th 2020. Available at: <https://www.cebm.net/covid-19/supporting-people-with-long-term-conditions-ltcs-during-national-emergencies/>
- 25) Qu X, Zhou XD. [Psychological intervention in oral patients in novel coronavirus pneumonia outbreak period] [Zhonghua Kou Qiang Yi Xue Za Zhi](#). 2020 Feb 22;55(0):E003. Article in Chinese. doi: 10.3760/cma.j.cn112144-20200213-00053. [Epub ahead of print]



iADH Covid-19 Fact Sheet by IADH is licensed under a

[Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](#).

Authors: Denise Faulks and Alison Dougall, in collaboration with Suzanne Krämer, Blanaid Daly and Pedro Diz Dios.

Comments, corrections, additions are welcome: vicechair@iadh.org